

# EPIDEMIOLOGY OF BLADDER CANCER

Debra T. Silverman, ScD, Patricia Hartge, ScD,  
Alan S. Morrison, MD, and Susan S. Devesa, PhD

Each year in the United States, approximately 49,000 persons develop cancer of the urinary bladder and 9700 die from the disease.<sup>182</sup> Bladder cancer accounts for 7% of all new cases of cancer among men and 3% of cases among women, as well as 2% of cancer deaths among men and 1% among women.

## DESCRIPTIVE FACTORS

### Histopathology and Anatomic Distribution

More than 98% of bladder cancers diagnosed in the United States are histologically confirmed (Devesa, unpublished data, and the Surveillance, Epidemiology, and End Results (SEER) program<sup>224</sup>). Most of these are transitional cell carcinomas (93%); 2% are squamous cell carcinomas, and 1% are adenocarcinomas. About 43% do not have a subsite within the bladder specified, and 13% arise in more than one subsite. Of bladder cancers for which a single subsite is specified, most occur on one of the bladder walls, with three times as many on the lateral walls (40%) as on the anterior (3%) or posterior (11%) walls combined. Less common subsites include the ureteric orifice (17%),

---

This article was adapted from Silverman DT, Morrison AS, Devesa SS: Bladder cancer. In Schotterfeld D, Fraumeni JF Jr (eds): *Cancer Epidemiology and Prevention*, ed 2. New York, Oxford University Press, in press

From the Epidemiology and Biostatistics Program (DTS, PH, SSD), Division of Cancer Etiology, National Cancer Institute, Bethesda, Maryland; and Department of Community Health (ASM), Brown University, Providence, Rhode Island

---

HEMATOLOGY/ONCOLOGY CLINICS OF NORTH AMERICA

VOLUME 6 • NUMBER 1 • FEBRUARY 1992

1

followed by the trigone, dome, and neck (13%, 9%, and 7%, respectively).

### Geographic Variation

Internationally, incidence rates of bladder cancer vary about tenfold.<sup>150</sup> High rates occur in western Europe and North America; relatively low rates are found in eastern Europe and several areas of Asia (Fig. 1). The geographic variation may reflect local practices in registration of "benign" tumors or "papillomas" as cancer, although rates reported by registries that include these categories are not consistently high compared with rates reported by other registries. Within the United States, bladder cancer incidence (Table 1) and mortality<sup>118, 166</sup> are higher in parts of the northeast and upper midwest and generally lower in the south.

In the United States, urban areas had higher mortality rates than rural areas, but these differences have diminished.<sup>10, 68</sup> Recent international incidence data indicate that urban excesses persist in many other countries.<sup>150</sup>


### Sex, Race, and Age

Cancer of the bladder occurs primarily among white men (Table 2). In most racial groups, the male:female rate ratio is at least 3:1, approaching 4:1 among whites. The incidence rate among black men is about 50% of that among whites. Rates are somewhat lower among Asian and Hispanic groups than among blacks, and very low among American Indians. Social class has little impact on the risk of bladder cancer and does not explain the black-white difference in incidence.<sup>48, 114, 125</sup>

Incidence and mortality rates rise sharply with age (Figs. 2 and 3); about two thirds of cases occur among persons 65 years of age and older. A higher incidence among whites compared with blacks occurs over the entire age range (see Fig. 2). A higher mortality in whites than in blacks is seen only among men 65 years of age and older (see Fig. 3). Among women, higher mortality in blacks than in whites is apparent at all ages except the oldest. Lifetime risk of developing bladder cancer is 2.8% for white men; for black men, white women, and black women, the risks are 0.9%, 1.0%, and 0.6%, respectively.<sup>179</sup>

### Stage of Disease

The stage of bladder cancer at diagnosis varies markedly by age, sex, and race, with young white women diagnosed with least advanced disease. The proportion localized at diagnosis declines with age; 82%



**Figure 1.** International variation in age-adjusted (world standard) bladder cancer incidence rates per 100,000 person-years by sex, 1978 to 1982. (Data from Muir C, Waterhouse J, Mack T, et al (eds): Cancer Incidence in Five Continents, vol. 5. IARC Publication Number 88. Lyon, International Agency for Research on Cancer, 1987.)

of patients aged 40 to 44 years present with local disease, whereas 61% of patients over 84 years do so (Devesa, unpublished data from the SEER program). The proportion localized is 73% among white men, 69% among white women, 59% among black men, and 49% among black women. The higher incidence among whites as compared with blacks is limited to localized cases, with blacks and whites having a similar risk of more advanced tumors.<sup>175</sup>

**Table 1. BLADDER CANCER INCIDENCE AMONG WHITES BY AREA AND SEX, US SEER PROGRAM, 1975-1985\***

Geographic area	Males		Females	
	Cases	Rate†	Cases	Rate†
Detroit	5127	33.5	1793	8.4
Hawaii	372	33.5	87	6.9
Connecticut	5153	32.7	1968	8.9
Seattle	3772	30.4	1322	8.0
San Francisco	3925	28.9	1536	7.9
Iowa	4686	28.3	1599	6.7
Atlanta	1312	28.3	477	6.7
Utah	1173	22.3	352	5.2
New Mexico	1064	19.9	348	5.3
All areas	26584	29.6	9482	7.6

\*Data from Devesa, unpublished data from the Surveillance, Epidemiology, and End Results (SEER) program.

†Rates per 100,000 person-years, age-adjusted by the direct method using the 1970 US population standard.

### Time Trends

From 1969-1971 to 1984-1986, incidence rose 22% to 38%, depending on race and sex. Increases in incidence were greater during the 1970s than during the 1980s and mortality fell 16% to 23% (Fig. 4).

The observed increases in incidence may be explained partly by changes in diagnostic practice. The proportion of bladder tumors classified as "carcinoma in situ" increased from less than 1% in 1969 to 1971 to more than 7% during the 1980s<sup>41</sup> (Devesa, unpublished data from the SEER program). Much of the observed rise in incidence

**Table 2. BLADDER CANCER INCIDENCE BY RACIAL/ETHNIC GROUP AND SEX, US SEER PROGRAM, 1975-1985\***

Racial/ethnic group	Males		Females	
	Cases	Rate†	Cases	Rate†
White	26584	29.6	9482	7.6
Black	1071	15.1	533	5.6
Chinese‡	157	13.9	48	3.9
Japanese‡	214	12.6	98	4.4
Hispanic§	178	11.3	61	3.3
Hawaiian	44	10.6	30	6.0
Filipino‡	72	5.9	21	3.0
American Indian§	10	3.5	1	0.4


\*Data from Devesa, unpublished data from the SEER program.

†Rates per 100,000 person-years, age-adjusted by the direct method using the 1970 US population standard.

‡Residents of San Francisco-Oakland and Hawaii.


§Residents of New Mexico.

||Hawaii only.




**Figure 2.** Age-specific incidence rates for bladder cancer in the US SEER program by race and sex, 1975 to 1985. (Data from the SEER program, unpublished data.)

appears to be a result of an increase in the incidence of localized bladder cancer (including in situ), which rose from 10.3 per 100,000 in 1975 to 1978 to 11.8 per 100,000 in 1982 to 1985 (Devesa, unpublished data from the SEER program). This increase was accompanied, however, by



**Figure 3.** Age-specific mortality rates for bladder cancer in the total United States by race and sex, 1975 to 1985. (Data from the National Center for Health Statistics.)




**Figure 4.** Trends in age-adjusted (1970 standard) bladder cancer incidence and mortality rates in the United States by race and sex, 1969 to 1986. Incidence data not available for 1972 to 1974. (Based on unpublished data from the five geographic areas (5GA) included in the 1969 to 1971 Third National Cancer Survey (TNCS), the SEER program, and the Connecticut Tumor Registry.<sup>49</sup>)

a decrease in the incidence of unstaged bladder cancer (1.7 to 0.7 per 100,000), suggesting that some of the apparent increase in localized disease was the result of a reduction in the frequency of unstaged cases. Incidence of regional- and distant-stage bladder cancer remained virtually constant.

### Survival

The 5-year relative survival of bladder cancer patients ranges from 87% for those diagnosed with localized disease to 9% for those with distant disease (Fig. 5). Whites with localized or regional disease have a better prognosis than blacks. White men with regional disease have a better prognosis than white women. Black women with localized disease have a worse prognosis than black men. Among cases with distant disease, survival varies little by race or sex. Overall black-white survival differences are only partly explained by differences in stage at diagnosis; racial disparities persist after adjustment for stage, histologic type, grade, and socioeconomic status.<sup>70, 161</sup>

Survival among patients with bladder cancer has increased more than 50% since the early 1960s.<sup>155</sup> In the period from 1974-1976 to 1980-1985, 5-year survival increased more among blacks (47.0% to 56.1%) than among whites (73.3% to 77.7%).



**Figure 5.** Five-year bladder cancer relative survival rates by stage, sex, and race. (Data from the National Cancer Institute, 1989.)

## RISK FACTORS

### Occupation

The study of occupational causes of bladder cancer gained momentum in the 1950s with the identification of bladder cancer hazards in the British dyestuffs and rubber industries.<sup>24, 25</sup> Numerous subsequent studies have identified approximately 40 high-risk occupations. Despite prodigious efforts, epidemiologists remain uncertain of the exact level of risks involved. Typically, a small number of workers are studied and their risk of bladder cancer is less than twice that expected. Further, many reported associations are not consistently found.<sup>187</sup> Strong evidence of increased risk is apparent for very few occupations: dye worker, aromatic amine manufacturing worker, leather worker, rubber worker, painter, truck driver, and aluminum worker.

### Dyestuffs Workers and Dye Users

In 1895, Rehn first suggested that men employed in the dyestuffs industry had increased risk of bladder cancer. In 1954, Case et al<sup>25</sup> showed that dyestuffs workers in England and Wales had a 10 to 50 fold increased risk of death from bladder cancer due to exposure to

two aromatic amines, 2-naphthylamine and benzidine. Exposure to aniline did not affect risk, and 1-naphthylamine apparently affected risk only because it was contaminated by 2-naphthylamine.

Dyestuffs workers in northern Italy<sup>14, 174</sup> experienced increased risk from exposure to 2-naphthylamine and benzidine. The ratios of observed to expected bladder cancer mortality were 13, 34, and 71 for dyestuffs workers employed 10 years or less, 11 to 20 years, and more than 20 years, respectively. Dyestuffs workers involved in fuchsin and safranin T manufacturing also experienced high mortality (observed/expected = 62.5), which may have been the result of exposure to two precursors, o-toluidine and 4,4'-methylene bis(2-methylaniline).

The mean time from start of exposure to death was 25 years, with a range of 12 to 41 years. Risk was greatest for workers who started before age 25 years (observed/expected = 200.0). Risk also decreased after exposure ceased, although no trend of decreasing risk with increasing time since last exposure was observed.

The increased risk among dyestuffs workers has also been observed in case-control studies,<sup>12, 109, 145, 153, 169, 171, 210</sup> with relative risks ranging from 1.7 to 8.8. Data from the United Kingdom indicate that bladder cancer risk among dyestuffs workers has been reduced since the introduction of protective measures and the subsequent banning of the industrial use of 2-naphthylamine and benzidine in 1950 and 1962, respectively.<sup>12, 145</sup>

Users of finished dyes also may have an increased risk of bladder cancer, but the evidence is not as persuasive as that for dyestuffs manufacturing workers. Kimono painters, many of whom ingest benzidine-based dyes by licking the brush, had seven times the expected rate of bladder cancer.<sup>223</sup> Coarse fishermen who use chrysoidine azo dyes to stain maggot bait have been reported to have an increased risk of bladder cancer (relative risk = 3.0 for fishermen who used bronze dyes for 5 or more years).<sup>192</sup> Canadian dyers of cloth were reported to have a relative risk of 4.6,<sup>171</sup> and British textile dyers with more than 20 years employment had a relative risk of 3.4,<sup>2</sup> but two other studies found no excess risk for dye users.<sup>23, 187</sup>

### *Aromatic Amine Manufacturing Workers*

Workers outside the dyestuffs industry have been exposed to 2-naphthylamine, benzidine, and a third bladder carcinogen, 4-aminobiphenyl, notably workers in chemical manufacturing.<sup>93</sup> A fourfold risk was observed among chemical workers in the United States exposed to 2-naphthylamine.<sup>177</sup> When excess bladder cancer was seen among workers involved in the commercial preparation of 4-aminobiphenyl, production ceased, averting widespread use of this aromatic amine.<sup>93</sup> In a cohort of workers at a benzidine manufacturing facility, an overall excess of bladder cancer cases was apparent (standardized incidence ratio [SIR] = 343).<sup>127</sup> Risk was greatest among those in the highest exposure category (SIR = 1303); little or no excess was observed for those in the low or medium exposure categories. Corresponding to the



introduction of preventive measures in the plant, a reduction in risk was observed for those first employed in 1950 or later compared with those first employed in 1945–49.

Two structural analogues of benzidine, MDA (4,4'-methylene-dianiline), and MBOCA (4,4'-methylene-bis[2-chloroaniline]), are carcinogenic in animals,<sup>178</sup> and possibly in humans as well. MDA, a curing agent for certain resins, was associated with a threefold elevation of proportional mortality from bladder cancer.<sup>178</sup> MBOCA, a curing agent used in the manufacture of rigid plastics, has been suggested as the exposure responsible for two noninvasive papillary tumors of the bladder in workers in a MBOCA production plant, although no invasive bladder tumors have been identified in the cohort.<sup>213</sup> Manufacturing of another dyestuffs intermediate, 4-chloro-o-toluidine (4-COT), has been associated with excess bladder cancer mortality in a cohort of chemical workers in Germany (relative risk = 72.7).<sup>193</sup>

### *Rubber Workers*

Antioxidants containing 2-naphthylamine were used in the rubber and electric-cable manufacturing industries in Great Britain until 1949.<sup>7, 163</sup> Case and Hosker<sup>24</sup> observed a twofold bladder cancer mortality risk, but only among British rubber workers employed before 1950.

Some studies of rubber workers in the United States<sup>26, 47, 140</sup> showed increased risk, but less than that seen among workers in Britain or Italy<sup>157</sup>; some studies found no excess risk.<sup>99, 109, 124, 145</sup> In the US rubber industry, workers were seldom exposed to 2-naphthylamine<sup>26</sup> but were exposed to another antioxidant, phenyl-β-naphthylamine (PBNA), which can be metabolized to 2-naphthylamine.<sup>93</sup>

### *Leather Workers*

Leather workers experienced an increased risk of bladder cancer in at least ten studies,<sup>8, 37, 45, 63, 77, 117, 145, 180, 210, 220</sup> but leather tanners in two studies did not.<sup>52, 197</sup> The relative risk has varied from 1.4 to 6.3, with "leather worker" defined differently among the studies. Increased risks have been seen for shoe makers and shoe repairers,<sup>63, 220</sup> for workers in leather products manufacturing<sup>45</sup> or, more broadly, for workers exposed to leather or leather products.<sup>37, 117, 145</sup>

Leather workers are exposed to leather dust and to dyes, their solvents, and unreacted intermediates.<sup>171</sup> The exposure responsible for the increased risk among leather workers is not known; however, Cole et al<sup>37</sup> observed that the excess was related to cutting, assembling, finishing, and related processes. Marrett et al<sup>117</sup> found risk to be slightly higher for workers with possible exposure to leather dust as compared with other types of leather exposure.

### *Painters*

Painters have had elevated bladder cancer risk in many studies,<sup>31, 37, 46, 69, 77, 99, 109, 116, 120, 133, 145, 152, 187, 220</sup> but not all.<sup>55, 145</sup> Typically, the overall

relative risks have been 1.2 to 1.5, and higher for those employed 20 years or more or before 1930 (3.0 to 4.1).<sup>99, 187</sup> Painters may be exposed to many known or suspected carcinogens in paints (e.g., benzidine, polychlorinated biphenyls, formaldehyde, and asbestos) and solvents (e.g., benzene, dioxane, and methylene chloride).<sup>133</sup>

#### *Drivers of Trucks and Other Motor Vehicles*

Excess risk of bladder cancer has been observed frequently among drivers of trucks, buses, or taxi cabs,<sup>8, 31, 46, 50, 80, 94, 99, 114, 131, 176, 183, 184, 194</sup> although one Swedish study found no elevation in risk for truck drivers.<sup>116</sup> Overall relative risks varied from 1.3 to 2.2, with relative risks for long-term drivers ranging from 2.2 to 12.0.<sup>31, 80, 99, 183, 184, 194</sup> In the largest study of bladder cancer among truck drivers, the trend in risk by duration of employment was most consistent for those first employed at least 50 years before observation<sup>184</sup> (Table 3). Although the etiologic agent has not been identified, one likely candidate is motor exhaust. Exhaust emissions contain polycyclic aromatic hydrocarbons (PAHs) and nitro-PAHs, which are highly mutagenic as well as carcinogenic in laboratory animals.<sup>184</sup>

#### *Aluminum Workers*

Wigle<sup>214</sup> noted an elevated incidence of bladder cancer among men in Quebec Province of Canada near the aluminum refining industry. The link was confirmed in two cohort studies of aluminum smelter workers<sup>65, 172</sup> and pinpointed to the Soderberg potrooms (relative risk = 2.4).<sup>199, 200</sup> Risk increased with increasing duration of employment in this department. Relative risks were 1.0 for less than 1 year, 1.9 for 1

**Table 3. BLADDER CANCER RISK AMONG TRUCK DRIVERS**

Duration of employment (yr)	Cases*	Controls*	Relative risk†
Never any motor exhaust-related occupation	1353	2724	1.0
<5	74	129	1.2
5-9	32	45	1.4
10-24	33	31	2.1
25+	22	19	2.2
(χ = 3.93; P < 0.0001)			

\*Numbers of cases and controls and relative risk according to duration of employment as a truck driver or deliveryman among those first employed at least 50 years before observation. The time of observation was the date of diagnosis for cases and the date of interview for controls. Males with unknown smoking history, duration of employment, or date started employment were excluded.

†Relative to a risk of 1.0 for males never employed in a motor exhaust-related occupation; adjusted for age and smoking.

‡Data from Silverman DT, Hoover RN, Mason TJ, et al: Motor exhaust-related occupations and bladder cancer. *Cancer Res* 46:2113-2116, 1986.

to 9 years, 3.0 for 10 to 19 years, 3.2 for 20 to 29 years, and 4.5 for 30 years or more.<sup>200</sup> Armstrong et al<sup>5</sup> used historical data on workplace exposures and found a tenfold risk gradient by exposure level (Table 4). The Soderberg electrolytic reduction process emits coal-tar pitch volatiles,<sup>200</sup> which may contain bladder carcinogens, such as aromatic amines (particularly 2-naphthylamine).<sup>5</sup>

### Other Occupations

Bladder cancer risk has been linked to employment as a machinist,<sup>187</sup> but not to a specific type of machine work. Machinists are exposed to mists from oils used as coolants and lubricants in metal machining processes,<sup>183, 207</sup> some of which contain potentially carcinogenic PAHs<sup>183</sup> and nitrosamines.<sup>56</sup>

Increased risk of bladder cancer has also been reported for many other occupational groups: metal workers, printers, chemical workers (other than those involved in manufacturing aromatic amines), hairdressers, dry cleaners, carpenters, construction workers, miners, gas workers, coke plant workers, auto mechanics, petroleum workers, railroad workers, textile workers, tailors, engineers, butchers, clerical workers, cooks and kitchen workers, food processing workers, electricians, gas station attendants, medical workers, pharmacists, glass processors, nursery workers, photographic workers, security guards and watchmen, welders, sailors, stationary fire fighters or furnace operators, stationary engineers, paper and pulp workers, roofers, gardeners, bootblacks, and asbestos workers.<sup>1, 119, 185, 187</sup> Findings for most of these occupations are not as persuasive as those discussed earlier and require corroboration.

The relation between occupation and bladder cancer risk is dynamic.<sup>187</sup> As bladder carcinogens are removed from the workplace and

**Table 4. RELATIVE RISKS PREDICTED FOLLOWING 40 YEARS OF EXPOSURE TO TAR VOLATILES\***

BSM			BaP		
Concentration (mg/m <sup>3</sup> )	Relative risk†	95% CI	Concentration (μg/m <sup>3</sup> )	Relative risk†	95% CI
1.0	8.1	3.8–17.4	10	10.2	4.6–21.8
0.5	4.5	2.40–9.2	5	5.6	2.8–11.4
0.2	2.42	1.56–4.3	2	2.84	1.72–5.2
0.1	1.71	1.28–2.64	1	1.92	1.36–2.15
0.05	1.35	1.14–1.82	0.5	1.46	1.18–2.04
0.02	1.14	1.06–1.33	0.2	1.18	1.07–1.42
0.01	1.07	1.03–1.16	0.1	1.09	1.04–1.21

BSM = benzene-soluble matter; BaP = benzo-a-pyrene; 95% CI = 95% confidence interval.

\*Assumes that a minimum of 10 years elapses before an effect of exposure occurs.

†Estimates of risk are relative to a risk of 1.0 for unexposed persons.

From Armstrong BG, Tremblay CG, Cyr D, et al: Estimating the relationship between exposure to tar volatiles and the incidence of bladder cancer in aluminum smelter workers. *Scand J Work Environ Health* 12:486–493, 1986; with permission.

new chemicals are introduced, changing worker exposures are generating shifts in "high-risk occupations." For example, risks among rubber and leather workers have diminished over time,<sup>117, 163</sup> whereas new high-risk occupations have emerged, such as truck driver and aluminum smelter worker.<sup>184, 214</sup> Thus, occupational bladder cancer continues to be a public health problem, with risks changing over time and from population to population.

## **Tobacco**

### **Cigarettes**

Cigarette smoking is accepted as a cause of human bladder cancer, as demonstrated in more than 30 case-control studies and in ten cohort studies.<sup>6, 16, 18, 32, 34, 42, 67, 75, 92, 94, 97, 135, 158, 170, 176</sup> Typical smokers have two to three times the risk of nonsmokers. Data from correlational studies also are consistent with a smoking-bladder cancer association. In the United States, bladder cancer mortality rates at the state level are highly correlated with per capita cigarette sales,<sup>61</sup> and birth cohort-specific patterns of bladder cancer incidence and mortality parallel the smoking patterns of those cohorts.<sup>3, 81</sup>

In nearly all studies, risk increases with increasing intensity of smoking (packs per day), with relative risk estimates for moderate to heavy smokers usually ranging from about 2.0 to 5.0 as compared with nonsmokers.<sup>6, 18, 32, 34, 42, 75, 92, 170, 176, 208</sup> Some studies report a plateau of risk at moderately high smoking levels; others show rising risk.<sup>6, 42, 75, 92, 176</sup> Longer duration of smoking also correlates with increased risk.<sup>6, 18, 32, 34, 42, 75, 92, 208</sup>

### **Cessation**

Cessation of cigarette smoking leads to an estimated 30% to 60% reduction in bladder cancer risk, as shown in many studies.<sup>92</sup> In some studies, former smokers who stopped smoking many years earlier had the same risk as nonsmokers.<sup>21, 42, 221</sup> Other studies, however, have shown that risk falls markedly within the first 2 to 4 years after stopping, but does not continue to decline thereafter and does not return to the level for nonsmokers.<sup>6, 18, 75, 208</sup> This reduction in risk within the first few years after quitting suggests that cigarette smoke contains agents that act at a late stage of bladder carcinogenesis.<sup>73</sup>

### **Filtration**

Persons who smoke unfiltered cigarettes exclusively appear to face a 35% to 50% higher risk of bladder cancer than those who smoke filtered cigarettes exclusively,<sup>75, 218</sup> yet switching to filtered cigarettes has not been shown to reduce the excess risk<sup>18, 75, 218</sup> (Table 5). This conundrum may mean that people who started smoking unfiltered

**Table 5. ESTIMATED RELATIVE RISKS OF BLADDER CANCER, ACCORDING TO USE OF FILTERED AND UNFILTERED CIGARETTES, AMONG CURRENT SMOKERS**

Filtered cigarettes/day	Unfiltered cigarettes/day			
	None*	1-19	20-39	>40
None		2.4 (1.3-4.5)	3.1 (1.7-5.6)	3.6 (1.8-6.9)
1-19	1.0	2.4 (1.4-4.1)	2.7 (1.3-5.5)	2.7 (0.8-8.5)
20-39	1.9 (1.1-3.3)	2.1 (1.2-3.7)	3.2 (1.9-5.5)	3.2 (1.5-6.7)
≥40	3.0 (1.4-6.5)	2.9 (1.2-7.0)	3.6 (2.0-6.6)	3.9 (2.1-7.1)
No. controls, cases				
None		87,56	172,40	61,57
1-19	102,29	165,122	35,28	8,6
20-39	90,48	100,68	328,321	26,26
≥40	24,21	16,15	71,79	73,85

\*Reference category is less than 20 filtered cigarettes per day, never smoked unfiltered. Estimates are adjusted for age, sex, race, and duration of smoking. 95% confidence intervals are shown in parentheses.

Data from Hartge P, Silverman D, Hoover R, et al: Changing cigarette habits and bladder cancer risk: A case-control study. J Natl Cancer Inst 78:1119-1125, 1987.

cigarettes start younger, take more puffs of smoke per cigarette, or inhale more deeply. Alternatively, changing from unfiltered to filtered cigarettes actually may exert almost no effect. Interview data may be too inaccurate to capture these subtle differences. If so, the observation that smokers of filtered cigarettes only have a lower risk than smokers of unfiltered cigarettes only needs further evaluation; the observed reduction in risk may have been a chance effect.

### *Inhalation*

Cigarette smokers who inhale deeply may have a greater risk than those who do not,<sup>18, 34, 38</sup> although some studies find no effect.<sup>75, 85, 113</sup> Morrison et al<sup>147</sup> observed 30% to 40% elevation of risk for male cigarette smokers who inhaled deeply as compared with those who inhaled somewhat or not at all.

### *Black versus Blond Tobacco*

Smokers of black tobacco have a risk of bladder cancer two to three times higher than the risk in smokers of blond tobacco.<sup>34, 43, 94, 208, 209</sup> Black tobacco has higher concentrations of aromatic amines, some of which are human bladder carcinogens, than does blond tobacco.<sup>208</sup> Blood levels of 4-aminobiphenyl hemoglobin adducts, as well as adducts

of several other aromatic amines, are higher for smokers of black than of blond tobacco.<sup>17</sup> In addition, the urine of smokers of black tobacco is more mutagenic than is the urine of smokers of blond tobacco.<sup>115, 136</sup>

### *Pipes, Cigars, and Smokeless Tobacco*

Most studies find pipe smokers at moderately elevated risk of bladder cancer (relative risks: 1.3 to 3.9),<sup>32, 73, 85, 97, 113, 137, 147, 189, 216, 220</sup> but some studies find no association.<sup>18, 38, 219</sup> A clear dose-response relationship is rarely seen, but pipe smokers who inhale deeply appear to be at greatest risk.<sup>73, 85</sup>

Weak and inconsistent relationships have been observed between bladder cancer risk and the other forms of tobacco use. Cigar smoking increases risk (relative risks: 1.3 to 2.5) in some studies,<sup>32, 73, 113, 137, 189</sup> but not in others.<sup>18, 38, 85, 97, 101, 147, 216, 219, 220</sup>

Most studies have reported that use of snuff or chewing tobacco does not increase risk of bladder cancer.<sup>18, 38, 73, 85, 137, 220</sup> Snuff users who never smoked cigarettes had excess risk in one study,<sup>189</sup> and users of chewing tobacco had excess risk in two studies.<sup>137, 189</sup>

## **Dietary Factors**

### *Coffee Drinking*

The hypothesis that coffee drinking increases bladder cancer risk was suggested by an unexpected finding in a population-based, case-control study conducted in Massachusetts (relative risk = 1.3 for men and 2.5 for women).<sup>36</sup> Nine subsequent studies indicated little or no overall association in either sex<sup>22, 30, 67, 98, 100, 148, 170</sup> (ref. 167: women only; ref. 189: men only). Eight studies were positive for men, but not for women<sup>14, 32, 33, 62, 74, 85, 130, 219</sup>; four studies were positive for women, but not for men<sup>134, 141, 171</sup> (ref. 188: women only); and four studies suggested an overall positive association without reporting sex-specific risks.<sup>94, 111, 135, 153</sup> In most of the studies reported as positive, however, the relative risk of bladder cancer in typical coffee drinkers compared with non-drinkers has been less than 2, and a regular dose-response relationship has been observed only infrequently,<sup>14, 32, 33, 94, 167, 219</sup> although risk was elevated among drinkers of large amounts of coffee in several studies.<sup>74, 98, 148, 170</sup>

The weakness and inconsistency of the observed associations indicate that if coffee is a bladder carcinogen, it is a weak one. Alternatively, associations between coffee drinking and bladder cancer could reflect residual confounding by cigarette smoking, a strong risk factor for bladder cancer and a strong correlate of coffee drinking.<sup>74, 143, 148</sup> Statistical adjustment is incomplete if smoking categories are too broad or if smoking habits are inaccurately recalled. This residual confounding by smoking can be avoided by evaluating the effect of coffee drinking in lifelong nonsmokers, but few studies have had

enough nonsmokers to estimate this risk with reasonable precision. Of these, some indicated no increased risk associated with coffee drinking,<sup>14, 85, 100, 148</sup> whereas others suggested an increased risk.<sup>30, 33, 74, 135, 170, 171, 189</sup> Of the positive studies that distinguished between men and women in examining the coffee drinking effect, one is positive in both men and women,<sup>170</sup> three are positive in men but not in women,<sup>30, 33, 74</sup> and one is positive in women but not in men.<sup>171</sup>

### *Artificial Sweeteners*

Artificial sweeteners were suggested as potential human bladder carcinogens by the results of animal experiments, particularly of rats exposed to high doses of saccharin in utero and weaned to a saccharin-containing diet.<sup>206</sup> Saccharin did not induce bladder cancer in rats or other animals fed saccharin only after birth.<sup>39</sup>

Extensive epidemiologic investigation has not substantiated a relationship between artificial sweeteners and bladder cancer. Diabetics, who often use artificial sweeteners, have typical bladder cancer mortality rates.<sup>4, 105</sup> Over time, the trend in bladder cancer mortality in England and Wales has not appeared related to saccharin consumption.<sup>3</sup> The Danish population born during World War II, a group with higher in utero saccharin exposure than previous birth cohorts, experienced no excess bladder cancer risk during the first 30 to 35 years of life.<sup>95</sup>

Of the numerous case-control studies measuring the relationship between artificial sweeteners and bladder cancer, most have been negative.<sup>22, 94, 96, 100, 106, 141, 146, 153, 167, 171, 188, 219, 222</sup> One study suggested an increased risk in men (relative risk = 1.6) and a decreased risk in women (relative risk = 0.6).<sup>84, 132</sup> In a large US population-based, case-control study, the relative risk for subjects who had ever used artificial sweeteners was 1.0.<sup>83</sup> A small elevation in risk was seen in very frequent users, but the dose-response pattern was irregular. A positive association was observed in two study subgroups: white male heavy smokers and white female nonsmokers with no known exposure to bladder carcinogens. The reason for these associations is uncertain, however.<sup>82, 212</sup>

It is difficult to separate the effects of saccharin and cyclamates in the United States and Canada because both substances were used extensively in both countries. In England and Japan, where primarily saccharin was used,<sup>147</sup> artificial sweeteners appear unrelated to bladder cancer risk.

The findings of nearly all studies indicate that the use of artificial sweeteners confers little or no excess risk of human bladder cancer. If, in fact, saccharin is a very weak carcinogen, such a low-level effect may not be detectable in epidemiologic studies.<sup>82</sup>

### *Alcohol Drinking*

Most studies of alcohol drinking and bladder cancer have shown no association.<sup>16, 22, 85, 100, 135, 153, 202, 219, 220</sup> Elevated risks related to con-

sumption of specific types of alcoholic beverages have been reported in a few studies,<sup>32, 94, 138, 141, 171, 189</sup> but these findings have not been consistent with respect to type of beverage or sex, and regular dose-response relationships have not been apparent. Thus, the positive findings probably reflect chance or residual confounding by smoking.

#### *Other Dietary Factors*

The role of dietary factors in human bladder carcinogenesis is unclear. Dietary supplements of natural and synthetic retinoids inhibit bladder carcinogenesis in laboratory animals,<sup>78</sup> but epidemiologic studies are inconsistent. Milk and other foods that contain vitamin A have been associated with decreased risk of bladder cancer in three case-control studies,<sup>111, 130, 189</sup> but two other case-control studies have not supported this relationship.<sup>171, 204</sup> The use of vitamin A supplements has been associated with decreased risk.<sup>195</sup> Serum levels of retinol, retinol binding protein, and carotenoids appear unrelated to risk.<sup>76, 159, 204</sup> Fruit and vegetable consumption has been associated with lower risk in some studies<sup>32, 111, 130, 135</sup> but not in others.<sup>196</sup>

Bladder cancer risk has been associated with higher intake of cholesterol,<sup>171</sup> with fatty meals,<sup>32</sup> with fried foods,<sup>195</sup> and with relatively high pork and beef consumption.<sup>196</sup> A nearly linear increasing trend in risk with decreasing serum levels of selenium was observed in a nested case-control study in Washington County, Maryland.<sup>76</sup>

### **Drugs**

#### *Analgesics*

Case reports first linked heavy consumption of phenacetin-containing analgesics to cancers of the renal pelvis, ureter, and bladder.<sup>90</sup> Fokkens<sup>60</sup> reported that Dutch subjects who had a lifetime consumption of at least 2 kg had a relative risk of 4.1 as compared with incidental users or nonusers. McCredie et al<sup>121, 122</sup> found a relative risk of 2.0 in Australian women 45 to 85 years of age who had a lifetime consumption of at least 1 kg. Piper et al<sup>168</sup> reported a relative risk of 6.5 in US women aged 20 to 44 years who had used phenacetin-containing compounds for at least 30 days in a year. Despite these fairly strong associations, a regular gradient in risk with increasing dose was demonstrated in only one study.<sup>122</sup> Further study will be limited because most western countries no longer allow phenacetin-containing analgesics to be sold.

Acetaminophen, an analgesic popular since the 1970s, showed no increased risk in two studies,<sup>121, 168</sup> but it may be too soon for any excess to be detectable.

#### *Cyclophosphamide and Chlornaphazine*

Cyclophosphamide, an alkylating agent used to treat both malignant and nonmalignant diseases since the early 1950s, has been linked



to risk of bladder cancer in many case reports and case series.<sup>91, 112</sup> Cyclophosphamide produces bladder tumors in both rats and mice.<sup>91</sup> Patients with non-Hodgkin's lymphoma treated with cyclophosphamide experienced a sevenfold risk of bladder cancer in a Danish study.<sup>165</sup> A report of bladder cancer risk among patients with non-Hodgkin's lymphoma who received chemotherapy, based on data from the SEER program and the Connecticut Tumor Registry, also was positive (observed/expected = 1.7).<sup>203</sup> Additional groups of patients, such as long-term survivors of breast cancer who were treated with cyclophosphamide as adjuvant chemotherapy, should be studied to measure more precisely the carcinogenic risk associated with use of this important antineoplastic drug.

In the 1960s, the antineoplastic drug chlornaphazine was linked to the development of bladder cancer.<sup>201</sup> Chlornaphazine is related chemically to 2-naphthylamine. This drug was never used in the United States and probably is not widely used elsewhere.<sup>89</sup>

## Urologic Conditions

### *Urinary Tract Infection*

Urinary bladder infection has been linked to risk of bladder cancer in several case-control studies<sup>51, 85, 102, 110, 167, 220</sup> but not all.<sup>107</sup> In the United States, Kantor et al<sup>102</sup> found an increased risk in both men and women; subjects with a history of at least three infections had a relative risk of 2.0, as compared with those with no infections. In addition, bladder infection was more strongly associated with squamous cell than with transitional cell cancer, a striking parallel to the relation between schistosomiasis and squamous cell bladder cancer. Because the dates of the bladder infections were not obtained in most of the reported studies, it is possible that the occurrence or diagnosis of infections may have been the consequence of early bladder cancer, rather than a cause of the disease.

### *Urinary Stasis*

If carcinogens are present in urine, urinary retention or stasis might increase the risk of developing bladder cancer by increasing the duration of contact of the carcinogens with the bladder mucosa.<sup>162</sup> Although urinary stasis has not been investigated directly as a risk factor, several findings are consistent with the hypothesis that stasis is related to risk. First, medical conditions that cause stasis, such as benign prostatic hypertrophy, have been associated with increased risk,<sup>51, 58, 139</sup> though it is uncertain whether these conditions preceded the diagnosis of bladder cancer. Second, infrequent micturition and high urine concentration, both of which increase urine contact with bladder epithelium, were more prevalent in high-risk areas of Israel than in low-risk areas.<sup>13</sup> Third, the upper hemisphere of the bladder (dome), which has less

contact with urine than the rest of the bladder, is a relatively infrequent site of bladder tumors.<sup>162</sup> Fourth, dogs exposed to 2-naphthylamine do not develop tumors in bladders that have not been in contact with urine.<sup>123</sup> Finally, urine itself appears to be a promoter of bladder carcinogenesis in the rat.<sup>160, 173</sup>

### *Schistosoma haematobium*

*S. haematobium* infection has been believed to increase risk of bladder cancer for 80 years,<sup>59</sup> and yet convincing epidemiologic data supporting this association are lacking. It is true that the proportional incidence of bladder cancer is high in countries with endemic schistosomiasis.<sup>198</sup> Also, squamous cell tumors are more common in endemic areas: in Egypt, more than 70% of bladder cancers are squamous cell<sup>198</sup>; in the United States, 2% are. On the other hand, bladder cancer incidence rates appear similar in areas of Africa where schistosomiasis is endemic, infrequent, or absent.<sup>164</sup>

In two studies, more bladder cancer patients had schistosome infection than did other hospital patients.<sup>64, 151</sup> In series of cases from South Africa and Zambia, *S. haematobium* ova were found in higher proportions of patients with squamous cell than with transitional cell tumors.<sup>9, 79</sup> Monkeys infected with *S. haematobium* develop bladder tumors,<sup>108</sup> but these are transitional cell. Hamsters infected with *S. haematobium* develop squamous metaplasia in the bladder.<sup>54</sup>

Schistosomiasis infection could predispose patients to bladder cancer if chronic inflammation by calcific ova and urinary retention caused by infection affects the absorption of carcinogens from the urine.<sup>27</sup> Alternatively, the urine of patients infected with *S. haematobium* or bacteria may harbor carcinogenic nitroso compounds,<sup>198</sup> or the schistosome antigen might depress the immunocompetence of infected patients.<sup>126</sup>

### Radiation

Ionizing radiation causes bladder cancer, but this exposure is so rare that it contributes very little to bladder cancer incidence in the general population. Women who received therapeutic pelvic radiation for dysfunctional uterine bleeding appear to have a two to four fold risk of bladder cancer.<sup>88, 211</sup> High-dose radiotherapy was associated with a fourfold risk of bladder cancer in a large, international study of cervical cancer patients treated with radiation.<sup>11</sup> Women under the age of 55 years when first treated had higher risks than older women. Risk increased with increasing dose to the bladder, and with time since exposure, with the relative risk reaching 8.7 for patients treated at least 20 years earlier.

Radioactive iodine (iodine-131) exposure has also been associated with elevated bladder cancer risk. A threefold risk was found among

women who had a thyroid uptake procedure with iodine-131.<sup>167</sup> A cohort of patients treated with high-dose iodine-131 for thyroid cancer also experienced excess risk.<sup>53</sup>

Follow-up of atomic bomb survivors in Hiroshima and Nagasaki revealed a dose-response relationship between radiation exposure and bladder cancer mortality.<sup>154</sup> Two groups of workers at nuclear installations in the United Kingdom had excess bladder cancer mortality,<sup>87, 191</sup> but no excess was seen in a group of US nuclear workers.<sup>66</sup>

### Drinking Water and Fluid Intake

Chlorination by-products in drinking water were first linked to bladder cancer risk by ecologic studies<sup>156</sup> and later by two case-control studies based on death certificates.<sup>40</sup>

In three investigations, detailed information was available on water quality and temporal aspects of exposure. These studies support the association between chlorination by-product levels in drinking water sources and bladder cancer risk. In Washington County, Maryland, residents supplied with chlorinated surface water had higher bladder cancer incidence rates than did those who consumed unchlorinated deep well water (relative risks were 1.8 and 1.6 for men and women, respectively).<sup>215</sup> In the NCI study conducted in ten areas of the United States, risk increased with level of intake of beverages made with tap water.<sup>19</sup> The gradient was restricted to subjects with at least 40 years of exposure to chlorinated surface water and was not observed among long-term consumers of nonchlorinated ground water. Among subjects whose residences were served by a chlorinated surface water source for at least 60 years, a relative risk of 2.0 was estimated for heavy consumers compared to low consumers of tap water. In a study in Massachusetts, residents of communities supplied with chlorinated drinking water experienced higher bladder cancer mortality than did those of communities exposed to water containing lower concentrations of chlorination by-products (mortality odds ratio = 1.6).<sup>225</sup>

A relation between exposure to high levels of arsenic in artesian well water and bladder cancer mortality has been suggested by surveys conducted in an endemic area of chronic arsenic toxicity, manifested by skin cancer and blackfoot disease in Taiwan.<sup>28, 29, 217</sup>

Total fluid intake also may be related to bladder cancer risk, but the results have been equivocal. Increased total fluid consumption has been associated with decreased risk,<sup>51</sup> with a positive trend in risk,<sup>19, 32, 98</sup> and with no excess risk.<sup>190, 220</sup>

### Hair Dyes

Use of hair dyes has been examined in relation to bladder cancer risk because hairdressers and barbers have been reported to be at

elevated risk.<sup>35</sup> In addition, findings from mutagenicity tests and animal experiments indicate that some compounds in hair dyes are mutagens and possible bladder carcinogens, and persons who dye their hair appear to excrete dye compounds in their urine.<sup>72</sup> Results of several epidemiologic studies, however, are negative.<sup>72</sup>

### Familial Occurrence

Familial predisposition to bladder cancer has been seen in clinical case reports and in a few case-control studies.<sup>20, 103, 167</sup> In the largest case-control study to date,<sup>103</sup> familial risks were especially high among those with environmental exposures, such as heavy cigarette smoking, suggesting genetic and environmental interactions.

### PREVENTIVE MEASURES

Avoidance of cigarette smoking is the most effective means available to prevent bladder cancer. In the United States, half of bladder cancer cases in men and one third in women are attributable to smoking, larger fractions than for any other single factor.<sup>92</sup> A further preventive measure would be curtailment of hazardous occupational exposures.

Screening offers some promise as a potential method to control bladder cancer. In general, a cancer screening program can be successful only if, first, an adequate screening test is available. Second, the prevalence of the detectable preclinical phase of the disease must be high. Third, a treatment must be available that is more effective for the screen-detected disease than for otherwise similar, routinely diagnosed disease.<sup>144</sup> It is uncertain whether all these conditions are met for bladder cancer.

Cytologic examination of the urine may be a satisfactory screening test, with a sensitivity of about 75%<sup>86, 104</sup> and a specificity above 95%,<sup>104</sup> possibly as high as 99.9%.<sup>86</sup> Furthermore, this test is inexpensive, convenient, and risk-free. Testing for asymptomatic hematuria also has been investigated as a method of screening for bladder cancer, but a single test is not sensitive enough, and repeated testing at short intervals<sup>128, 129</sup> would yield too many false positives.

Too few preclinical bladder cancers are prevalent in the general population for large-scale screening programs to be rewarding.<sup>142</sup> On the other hand, certain occupational groups probably have high enough prevalence to justify screening as a bladder cancer control measure.<sup>104</sup>

The natural history of preclinical bladder cancer is poorly understood. Screen-detected cases of most types of cancer typically have better survival than routinely diagnosed cases, even if screening confers no benefit in terms of reduced mortality from the disease.<sup>144</sup> For bladder cancer, the opposite pattern may obtain, with worse survival in screen-detected cases.<sup>181</sup> Screening for bladder cancer seems to detect prefer-

entially cases with prognostically unfavorable histologic appearance.<sup>57, 104</sup> Neither of these findings, by themselves, would negate the possible value of screening. However, they suggest that the gains may be small, and that particular care is needed in designing and evaluating bladder cancer screening programs.<sup>86</sup>

Lastly, the effectiveness of the treatment of bladder cancer is uncertain. No data are available on the relative value of treating bladder cancer "early," as detected by screening, or "late," as it comes to medical attention as a result of symptoms. In summary, research on screening probably should be focused on intervention trials among high-risk groups to determine whether urine cytology, alone or in combination with other screening methods, leads to a reduction in mortality from bladder cancer.<sup>205</sup>

## SUMMARY

Approximately 49,000 persons in the United States develop bladder cancer each year, and about 9700 die of it. White men face a lifetime risk of almost 3%; white women and black men face a risk of about 1%, and black women, about 0.5%.

Cigarette smoking is accepted widely as a cause of bladder cancer. Smoking accounts for about half of bladder cancer diagnosed among men and about one third of that among women. Moderate to heavy smokers typically show a two to five fold risk of bladder cancer, compared with persons who never smoked. When cigarette smokers quit smoking, their bladder cancer risk falls measurably within 2 to 4 years, but probably does not continue to decline with increasing years since quitting and does not appear to return to the baseline level of nonsmokers.

Occupational exposure to certain aromatic amines causes human bladder cancer. Clear evidence of bladder cancer risk also is apparent for a small number of occupational groups: dye workers, rubber workers, leather workers, painters, truck drivers, and aluminum workers. Many other occupational groups have been reported to have increased bladder cancer risk, but evidence for these is not as strong.

Coffee drinking has been studied extensively as a potential risk factor, but the inconsistency of the observed associations suggests that the relationship is either quite weak, noncausal, or dependent in a complex way on unmeasured factors. Artificial sweeteners confer little or no excess bladder cancer risk. Alcohol consumption apparently does not affect risk either. Consumption of fruits, vegetables, and foods high in vitamin A have been suggested as possible protective factors; consumption of high-fat foods, pork, and beef have been suggested as possible risk factors. Further epidemiologic research is needed to elucidate the role of diet in human bladder carcinogenesis.

Less common risk factors for bladder cancer include ionizing radiation, cyclophosphamide use, and abuse of phenacetin-containing

analgesics. Schistosomiasis infection may contribute substantially to the bladder cancer burden in Egypt and elsewhere, though not in the United States. Other urinary tract infections that are common worldwide may increase risk, but additional research is necessary to confirm that the infections preceded the occurrence of the bladder cancer.

Long-term consumption of drinking water containing chlorination by-products, total fluid volume, and urinary stasis have all been suggested to influence risk. Because these exposures are difficult to measure and to separate from one another, a clear conclusion regarding them cannot be drawn at this time.

Large-scale screening of the general population for early bladder cancer is not feasible yet. An intervention trial is needed to assess whether screening by urine cytology, alone or in combination with other methods, reduces mortality in a group at high risk for bladder cancer.

## References

1. Alderson M: Occupational Cancer. London, Butterworth and Co, 1986
2. Antony HM: Industrial exposure in patients with carcinoma of the bladder. *J Soc Occup Med* 24:110, 1974
3. Armstrong B, Doll R: Bladder cancer mortality in England and Wales in relation to cigarette smoking and saccharin consumption. *Br J Prev Soc Med* 28:233-240, 1974
4. Armstrong B, Doll R: Bladder cancer mortality in diabetics in relation to saccharin consumption and smoking habits. *Br J Prev Soc Med* 29:73-81, 1975
5. Armstrong BG, Tremblay CG, Cyr D, et al: Estimating the relationship between exposure to tar volatiles and the incidence of bladder cancer in aluminum smelter workers. *Scand J Work Environ Health* 12:486-493, 1986
6. Augustine A, Hebert JR, Kabat GC, et al: Bladder cancer in relation to cigarette smoking. *Cancer Res* 48:4405-4408, 1988
7. Baus Subcommittee on Industrial Bladder Cancer: Occupational bladder cancer: A guide for clinicians. *Br J Urol* 61:183-191, 1988
8. Baxter PJ, McDowall ME: Occupation and cancer in London: An investigation into nasal and bladder cancer using the Cancer Atlas. *Br J Ind Med* 43:44-49, 1986
9. Bhagwande SB: Schistosomiasis and carcinoma of the bladder in Zambia. *S Afr Med J* 50:1616-1620, 1976
10. Blot WJ, Fraumeni JF Jr: Geographic patterns of bladder cancer in the United States. *J Natl Cancer Inst* 61:1017-1023, 1978
11. Boice JD, Engholm G, Kleinerman RA, et al: Radiation dose and second cancer risk in patients treated for cancer of the cervix. *Radiat Res* 116:3-55, 1988
12. Boyko RW, Cartwright RA, Glashan RW: Bladder cancer in dye manufacturing workers. *J Occup Med* 27:799-803, 1985
13. Braver DJ, Modan M, Chetrit A, et al: Drinking, micturition habits, and urine concentration as potential risk factors in urinary bladder cancer. *J Natl Cancer Inst* 78:437-440, 1987
14. Bross ID, Tidings J: Another look at coffee drinking and cancer of the urinary bladder. *Prev Med* 2:445-451, 1973
15. Brown RR, Price JM, Friedell GH, et al: Tryptophan metabolism in patients with bladder cancer: Geographical differences. *J Natl Cancer Inst* 43:295-301, 1969
16. Brownson RC, Chang JC, Davis JR: Occupation, smoking, and alcohol in the epidemiology of bladder cancer. *Am J Publ Health* 77:1298-1300, 1987
17. Bryant MS, Vineis P, Skipper PL, et al: Hemoglobin adducts of aromatic amines. Associations with smoking status and type of tobacco. *Proc Natl Acad Sci USA* 85:9788-9791, 1988

18. Burch JD, Rohan TE, Howe GR, et al: Risk of bladder cancer by source and type of tobacco exposure: A case-control study. *Int J Cancer* 44:622-628, 1989
19. Cantor KP, Hoover R, Hartge P, et al: Bladder cancer, drinking water source, and tap water consumption: A case-control study. *J Natl Cancer Inst* 79:1269-1279, 1987
20. Cartwright RA: Genetic association with bladder cancer. *Br Med J* 2:798, 1979
21. Cartwright RA, Adib R, Appleyard I, et al: Cigarette smoking and bladder cancer: An epidemiological inquiry in West Yorkshire. *J Epidemiol Commun Health* 37:256-263, 1983
22. Cartwright RA, Adib R, Glashan R, et al: The epidemiology of bladder cancer in West Yorkshire. A preliminary report on non-occupational aetiologies. *Carcinogenesis* 2:343-347, 1981
23. Cartwright RA, Bernard SM, Glashan RW, et al: Bladder cancer amongst dye users. *Lancet* 2:1073-1074, 1979
24. Case RAM, Hosker ME: Tumour of the urinary bladder as an occupational disease in the rubber industry in England and Wales. *Br J Prev Soc Med* 8:39-50, 1954
25. Case RAM, Hosker ME, McDonald DB, et al: Tumours of the urinary bladder in workmen engaged in the manufacture and use of certain dyestuff intermediates in the British chemical industry. *Br J Ind Med* 11:75-104, 1954
26. Checkoway H, Smith AH, McMichael AJ, et al: A case-control study of bladder cancer in the United States rubber and tire industry. *Br J Ind Med* 38:240-246, 1981
27. Cheever AW: Schistosomiasis and neoplasia. *J Natl Cancer Inst* 61:13-18, 1978
28. Chen C-J, Chuang Y-C, Lin T-M, et al: Malignant neoplasms among residents of a blackfoot disease-endemic area in Taiwan: High-arsenic artesian well water and cancers. *Cancer Res* 45:5895-5899, 1985
29. Chen C-J, Chuang Y-C, You S-L, et al: A retrospective study on malignant neoplasms of bladder, lung and liver in blackfoot disease endemic area in Taiwan. *Br J Cancer* 53:399-405, 1986
30. Ciccone G, Vineis P: Coffee drinking and bladder cancer. *Cancer Lett* 41:45-52, 1988
31. Claude JC, Frentzel-Beyme RR, Kunze E: Occupation and risk of cancer of the lower urinary tract among men. A case-control study. *Int J Cancer* 41:371-379, 1988
32. Claude J, Kunze E, Frentzel-Beyme R, et al: Life-style and occupational risk factors in cancer of the lower urinary tract. *Am J Epidemiol* 124:578-589, 1986
33. Clavel J, Cordier S: Coffee consumption and bladder cancer risk. *Int J Cancer* 47:207-212, 1991
34. Clavel J, Cordier S, Boccon-Gibod L, et al: Tobacco and bladder cancer in males: Increased risk for inhalers and smokers of black tobacco. *Int J Cancer* 44:605-610, 1989
35. Clemmesen J: Epidemiological studies into the possible carcinogenicity of hair dyes. *Mutat Res* 87:65-79, 1981
36. Cole P: Coffee drinking and cancer of the lower urinary tract. *Lancet* 1:1335-1337, 1971
37. Cole P, Hoover R, Friedell G: Occupation and cancer of the lower urinary tract. *Cancer* 29:1250-1260, 1972
38. Cole P, Monson RR, Haning H, et al: Smoking and cancer of the lower urinary tract. *N Engl J Med* 284:129-134, 1971
39. Council on Scientific Affairs: Saccharin review of safety issues. *JAMA* 254:2622-2624, 1985
40. Crump KS, Guess HA: Drinking water and cancer: Review of recent epidemiological findings and assessment of risks. *Ann Rev Publ Health* 3:339-357, 1982
41. Cutler SJ, Young JL Jr (eds): Third National Cancer Survey: Incidence Data. *Natl Cancer Inst Monogr* 41:1-454, 1975
42. D'Avanzo B, Negri E, La Vecchia C, et al: Cigarette smoking and bladder cancer. *Eur J Cancer* 26:714-718, 1990
43. De Stefani E, Correa P, Fierro L, et al: Black tobacco, mate, and bladder cancer. *Cancer* 67:536-540, 1991
44. Decarli A, Peto J, Piolatto G, et al: Bladder cancer mortality of workers exposed to aromatic amines: Analysis of models of carcinogenesis. *Br J Cancer* 51:707-712, 1985
45. Decoufle P: Cancer risks associated with employment in the leather and leather products industry. *Arch Environ Health* 34:33-37, 1979

46. Decoufle P, Stanislawczyk K, Houten L, et al: A Retrospective Survey of Cancer in Relation to Occupation. DHEW Publ No. (NIOSH) 77-178. Washington, US Govt Printing Office, 1977
47. Delzell E, Monson RR: Mortality among rubber workers. III. Cause-specific mortality, 1940-1978. *J Occup Med* 23:677-684, 1981
48. Devesa SS: The Association of Cancer Incidence with Income and Education. Unpublished doctoral thesis, Johns Hopkins University, Baltimore, 1979
49. Devesa SS, Silverman DT, Young JL Jr, et al: Cancer incidence and mortality trends among whites in the United States, 1947-84. *J Natl Cancer Inst* 79:701-770, 1987
50. Dubrow R, Wegman DH: Cancer and occupation in Massachusetts: A death certificate study. *Am J Ind Med* 6:207-230, 1984
51. Dunham LJ, Rabson AS, Stewart HL, et al: Rates, interview, and pathology study of cancer of the urinary bladder in New Orleans, Louisiana. *J Natl Cancer Inst* 41:683-709, 1968
52. Edling C, Kling H, Flodin U, et al: Cancer mortality among leather tanners. *Br J Ind Med* 43:494-496, 1986
53. Edmonds CJ, Smith T: The long-term hazards of the treatment of thyroid cancer with radioiodine. *Br J Radiol* 59:45-51, 1986
54. El-Morsi B, Sherif M, El-Raziki ES: Experimental bilharzial squamous metaplasia of the urinary bladder in hamsters. *Eur J Cancer* 11:199-201, 1974
55. Englund A: Cancer incidence among painters and some allied trades. *J Toxicol Environ Health* 6:1267-1273, 1980
56. Fan TY, Morrison J, Rounbehler DP, et al: *N*-Nitrosodiethanolamine in synthetic cutting fluids: A part-per-hundred impurity. *Science* 196:70-71, 1977
57. Farrow GM: Pathologist's role in bladder cancer. *Semin Oncol* 6:198-206, 1979
58. Fellows GJ: The association between vesical carcinoma and urinary obstruction. *Eur Urol* 4:187-188, 1978
59. Ferguson AR: Associated bilharziosis and primary malignant disease of the urinary bladder with observations in a series of forty cases. *J Pathol Bacteriol* 16:76-94, 1911
60. Fokkens W: Phenacetin abuse related to bladder cancer. *Environ Res* 20:192-198, 1979
61. Fraumeni JF Jr: Cigarette smoking and cancers of the urinary tract: Geographic variation in the United States. *J Natl Cancer Inst* 41:1205-1211, 1968
62. Fraumeni JF Jr, Scotto J, Dunham LF: Coffee drinking and bladder cancer. *Lancet* 2:1204, 1971
63. Garabrant DH, Wegman DH: Cancer mortality among shoe and leather workers in Massachusetts. *Am J Ind Med* 5:303-314, 1984
64. Gelfand M, Weinberg RW, Castle WM: Relation between carcinoma of the bladder and infestation with *Schistosoma haematobium*. *Lancet* 1:1249-1251, 1967
65. Gibbs GW: Mortality of aluminum reduction plant workers, 1950 through 1957. *J Occup Med* 27:761-770, 1985
66. Gilbert ES, Fry SA, Wiggs LD, et al: Analyses of combined mortality data on workers at the Hanford Site, Oak Ridge National Laboratory, and Rocky Flats Nuclear Weapons Plant. *Radiat Res* 120:19-35, 1989
67. Gonzales CA, Lopez-Abente G, Errezola M, et al: Occupation, tobacco use, coffee, and bladder cancer in the county of Mataro (Spain). *Cancer* 55:2031-2034, 1985
68. Greenberg MR: Urbanization and Cancer Mortality: The United States Experience, 1950-1975. Monographs in Epidemiology and Biostatistics, vol 4. New York, Oxford University Press, 1983
69. Guberan E, Usel M, Raymond L, et al: Disability, mortality, and incidence of cancer among Geneva painters and electricians: A historical prospective study. *Br J Ind Med* 46:16-23, 1989
70. Hankey BF, Myers MH: Black/white differences in bladder cancer survival. *J Chron Dis* 40:65-73, 1987
71. Hartge P, Harvey EB, Linehan WM, et al: Explaining the male excess in bladder cancer risk. *J Natl Cancer Inst* 82:1636-1640, 1990
72. Hartge P, Hoover R, Altman R, et al: Use of hair dyes and risk of bladder cancer. *Cancer Res* 42:4784-4787, 1982
73. Hartge P, Hoover R, Kantor A: Bladder cancer risk and pipes, cigars and smokeless tobacco. *Cancer* 55:901-906, 1985



74. Hartge P, Hoover R, West DW, Lyon JL: Coffee drinking and risk of bladder cancer. *J Natl Cancer Inst* 70:1021-1026, 1983
75. Hartge P, Silverman D, Hoover R, et al: Changing cigarette habits and bladder cancer risk: A case-control study. *J Natl Cancer Inst* 78:1119-1125, 1987
76. Helzlsouer KJ, Comstock GW, Morris JS: Selenium, lycopene,  $\alpha$ -tocopherol,  $\beta$ -carotene, retinol, and subsequent bladder cancer. *Cancer Res* 49:6144-6148, 1989
77. Henry SA, Kennaway NM, Kennaway EL: The incidence of cancer of the bladder and prostate in certain occupations. *J Hyg* 31:125-137, 1931
78. Hicks RM: The scientific basis for regarding vitamin A and its analogues as anti-carcinogenic agents. *Proc Nutr Soc* 42:83-93, 1983
79. Hinder RA, Schmaman A: Bilharziasis and squamous carcinoma of the bladder. *S Afr Med J* 43:617-618, 1969
80. Hoar SK, Hoover R: Truck driving and bladder cancer mortality in rural New England. *J Natl Cancer Inst* 74:771-774, 1985
81. Hoover R, Cole P: Population trends in cigarette smoking and bladder cancer. *Am J Epidemiol* 94:409-418, 1971
82. Hoover R, Hartge P: Non-nutritive sweeteners and bladder cancer. *Am J Publ Health* 72:382-383, 1982
83. Hoover RN, Strasser PH, Child M, et al: Artificial sweeteners and human bladder cancer. *Lancet* 1:837-840, 1980
84. Howe GR, Burch JD, Miller AB, et al: Artificial sweeteners and human bladder cancer. *Lancet* 2:578-581, 1977
85. Howe GR, Burch JD, Miller AB, et al: Tobacco use, occupation, coffee, various nutrients, and bladder cancer. *J Natl Cancer Inst* 64:701-713, 1980
86. Hutchison GB: Summary. State-of-the-Art Conference on Bladder Cancer Screening. Washington, DC, 1977
87. Inskip H, Beral V, Fraser P, et al: Further assessment of the effects of occupational radiation exposure in the United Kingdom Atomic Energy Authority mortality study. *Br J Ind Med* 44:149-160, 1987
88. Inskip PD, Monson RR, Wagoner JK, et al: Cancer mortality following radiotherapy for uterine bleeding. *Radiat Res* 123:331-344, 1990
89. International Agency for Research on Cancer: Evaluation of Carcinogenic Risk of Chemicals to Man. Some Aromatic Amines, Hydrazine and Related Substances, N-nitroso Compounds and Miscellaneous Alkylating Agents, vol 4. Lyon, International Agency for Research on Cancer, 1974
90. International Agency for Research on Cancer: Evaluation of the Carcinogenic Risk of Chemicals to Humans. Some Pharmaceutical Drugs, vol 24. Lyon, International Agency for Research on Cancer, 1980
91. International Agency for Research on Cancer: Evaluation of the Carcinogenic Risk of Chemicals to Humans. Some Antineoplastic and Immunosuppressive Agents, vol 26. Lyon, International Agency for Research on Cancer, 1981
92. International Agency for Research on Cancer: Evaluation of the Carcinogenic Risk of Chemicals to Humans. Tobacco Smoking, vol 38. Lyon, International Agency for Research on Cancer, 1986
93. International Agency for Research on Cancer: Overall Evaluations of Carcinogenicity. An Updating of IARC Monographs Volumes 1 to 42, Suppl 7. Lyon, International Agency for Research on Cancer, 1987
94. Iscovich J, Castelletto R, Esteve J, et al: Tobacco smoking, occupational exposure and bladder cancer in Argentina. *Int J Cancer* 40:734-740, 1987
95. Jensen OM, Kamby C: Intra-uterine exposure to saccharine and risk of bladder cancer in man. *Int J Cancer* 29:507-509, 1982
96. Jensen OM, Knudsen JB, Sorensen BL, et al: Artificial sweeteners and absence of bladder cancer risk in Copenhagen. *Int J Cancer* 32:577-582, 1983
97. Jensen OM, Wahrendorf J, Blettner M, et al: The Copenhagen case-control study of bladder cancer: Role of smoking in invasive and non-invasive bladder tumours. *J Epidemiol Commun Health* 41:30-36, 1987
98. Jensen OM, Wahrendorf J, Knudsen JB, et al: The Copenhagen case-control study of bladder cancer. II. Effect of coffee and other beverages. *Int J Cancer* 37:651-657, 1986

99. Jensen OM, Wahrendorf J, Knudsen JB, et al: The Copenhagen case-referent study on bladder cancer. Risks among drivers, painters and certain other occupations. *Scand J Work Environ Health* 13:129-134, 1987
100. Kabat GC, Dieck GS, Wynder EL: Bladder cancer in nonsmokers. *Cancer* 57:362-367, 1986
101. Kahn HA: The Dorn study of smoking and mortality among U.S. veterans. Report on eight and one-half years of observation. In Haenszel W (ed): *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Natl Cancer Inst Monogr 19. Washington, US Govt Printing Office, 1966, pp 1-125
102. Kantor AF, Hartge P, Hoover RN, et al: Urinary tract infection and risk of bladder cancer. *Am J Epidemiol* 119:510-515, 1984
103. Kantor AF, Hartge P, Hoover RN, et al: Familial and environmental interactions in bladder cancer risk. *Int J Cancer* 35:703-706, 1985
104. Kern WH: Screening tests for bladder cancer. In Miller AB (ed): *Screening for Cancer*. New York, Academic Press, 1985, p 121
105. Kessler II: Cancer mortality among diabetics. *J Natl Cancer Inst* 44:673-686, 1970
106. Kessler II, Clark JP: Saccharin, cyclamate, and human bladder cancer. *JAMA* 240:349-355, 1978
107. Kjaer SK, Knudsen JB, Sorensen BL, et al: The Copenhagen case-control study of bladder cancer. *Acta Oncol* 28:631-636, 1989
108. Kuntz RE, Cheever AW, Myers BJ: Proliferative epithelial lesions of the urinary bladder of nonhuman primates infected with *Schistosoma haematobium*. *J Natl Cancer Inst* 48:223-235, 1972
109. La Vecchia C, Negri E, D'Avanzo B, et al: Occupation and the risk of bladder cancer. *Int J Epidemiol* 19:264-268, 1990
110. La Vecchia C, Negri E, D'Avanzo B, et al: Genital and urinary tract diseases and bladder cancer. *Cancer Res* 51:629-631, 1991
111. La Vecchia C, Negri E, Decarli A, et al: Dietary factors in the risk of bladder cancer. *Nutr Cancer* 12:93-101, 1989
112. Levine LA, Richie JP: Urological complications of cyclophosphamide. *J Urol* 141:1063-1069, 1989
113. Lockwood K: On the etiology of bladder tumors in København-Frederiksberg: An inquiry of 369 patients and 369 controls. *Acta Pathol Microbiol Scand* 51 (Suppl 145):1-166, 1961
114. Logan WPD: *Cancer Mortality by Occupation and Social Class, 1851-1971*. IARC Publ No. 36. Lyon, International Agency for Research on Cancer, 1982.
115. Malaveille C, Vineis P, Esteve J, et al: Levels of mutagens in the urine of smokers of black and blond tobacco correlate with their risk of bladder cancer. *Carcinogenesis* 10:577-586, 1989
116. Walker HSR, McLaughlin JK, Silverman DT, et al: Occupational risks for bladder cancer among men in Sweden. *Cancer Res* 47:6763-6766, 1987
117. Marrett LD, Hartge P, Meigs JW: Bladder cancer and occupational exposure to leather. *Br J Ind Med* 43:96-100, 1986
118. Mason TJ, McKay F, Hoover R, et al: *Atlas of Cancer Mortality for US Counties: 1950-1969*. DHEW Publ No. (NIH) 75-780. Washington, US Govt Printing Office, 1975
119. Matanoski GM, Elliot EA: Bladder cancer epidemiology. *Epidemiol Rev* 3:203-229, 1981
120. Matanoski GM, Stockwell HG, Diamond EL, et al: A cohort mortality study of painters and allied tradesmen. *Scand J Work Environ Health* 12:16-21, 1986
121. McCredie M, Stewart JH: Does paracetamol cause urothelial cancer or renal papillary necrosis? *Nephron* 49:296-300, 1988
122. McCredie M, Stewart JH, Ford JM, et al: Phenacetin-containing analgesics and cancer of the bladder or renal pelvis in women. *Br J Urol* 55:220-224, 1983
123. McDonald DF, Lund RR: The role of the urine in vesical neoplasm. 1. Experimental confirmation of the urogenous theory of pathogenesis. *J Urol* 71:560-570, 1954
124. McMichael AJ, Andjelkovich DA, Tyroler HA: Cancer mortality among rubber workers. *Ann NY Acad Sci* 271:125-137, 1976
125. McWhorter WP, Schatzkin AG, Horm JW, et al: Contribution of socioeconomic status to black/white differences in cancer incidence. *Cancer* 63:982-987, 1989

126. Mee AD: Aetiological aspects of bladder cancer in urinary bilharzia. *Saudi Med J* 3:123-128, 1982
127. Meigs JW, Marrett LD, Ulrich FU, et al: Bladder tumor incidence among workers exposed to benzidine: A thirty-year follow-up. *J Natl Cancer Inst* 76:1-8, 1986
128. Messing EM, Young TB, Hunt VB, et al: The significance of asymptomatic micro-hematuria in men 50 or more years old: Findings of a home screening study using urinary dipsticks. *J Urol* 137:919-922, 1987
129. Messing EM, Young TB, Hunt VB, et al: Urinary tract cancers found by homes using hematuria dipsticks in healthy men over 50 years of age. *Cancer* 64:2361-2367, 1989
130. Mettlin C, Graham S: Dietary risk factors in human bladder cancer. *Am J Epidemiol* 110:253-263, 1979
131. Milham S Jr: Occupational mortality in Washington State, 1950-1971. DHEW Publ No. (NIOSH) 76-175-C. Washington, US Govt Printing Office, 1976
132. Miller AB, Howe G: Artificial sweeteners and bladder cancer. *Lancet* 2:1221-1222, 1977
133. Miller BA, Silverman DT, Hoover RN, et al: Cancer risk among artistic painters. *Am J Ind Med* 9:281-287, 1986
134. Miller CT, Neutel CI, Nair RC, et al: Relative importance of risk factors in bladder carcinogenesis. *J Chron Dis* 31:51-56, 1978
135. Mills PK, Beeson L, Phillips RL, et al: Bladder cancer in a low risk population: Results from the Adventist Health Study. *Am J Epidemiol* 133:230-239, 1991
136. Mohtashamipur E, Norpoth K, Lieder F: Urinary excretion of mutagens in smokers of cigarettes with various tar and nicotine yields, black tobacco, and cigars. *Cancer Lett* 34:103-112, 1987
137. Mommsen S, Aagaard J: Tobacco as a risk factor in bladder cancer. *Carcinogenesis* 4:335-338, 1983
138. Mommsen S, Aagaard J, Sell A: An epidemiological case-control study of bladder cancer in males from a predominantly rural district. *Eur J Cancer Clin Oncol* 18:1205-1210, 1982
139. Mommsen S, Sell A: Prostatic hypertrophy and venereal disease as possible risk factors in the development of bladder cancer. *Urol Res* 11:49-52, 1983
140. Monson RR, Nakano KK: Mortality among rubber workers. I. White male union employees in Akron, Ohio. *Am J Epidemiol* 103:284-296, 1976
141. Morgan RW, Jain MG: Bladder cancer: Smoking, beverages and artificial sweeteners. *Can Med Assoc J* 111:1067-1070, 1974
142. Morrison AS: The public health value of using epidemiologic information to identify high risk groups for bladder cancer screening. *Semin Oncol* 6:184-188, 1979
143. Morrison AS: Control of cigarette smoking in evaluating the association of coffee drinking and bladder cancer. In MacMahon B, Sugimura T (eds): *Coffee and Health*, Banbury Report 17. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory, 1984, pp 127-136
144. Morrison AS: *Screening in Chronic Disease*. New York, Oxford University Press, 1985
145. Morrison AS, Ahlbom A, Verhoek WG, et al: Occupation and bladder cancer in Boston, USA, Manchester, UK, and Nagoya, Japan. *J Epidemiol Commun Health* 39:294-300, 1985
146. Morrison AS, Buring JE: Artificial sweeteners and cancer of the lower urinary tract. *N Engl J Med* 302:537-541, 1980
147. Morrison AS, Buring JE, Verhoek WG, et al: An international study of smoking and bladder cancer. *J Urol* 131:650-654, 1984
148. Morrison AS, Buring JE, Verhoek WG, et al: Coffee drinking and cancer of the lower urinary tract. *J Natl Cancer Inst* 68:91-94, 1982
149. Morrison AS, Verhoek WG, Leck I, et al: Artificial sweeteners and bladder cancer in Manchester, UK, and Nagoya, Japan. *Br J Cancer* 45:332-336, 1982
150. Muir C, Waterhouse J, Mack T, et al (eds): *Cancer Incidence in Five Continents*, vol V. IARC Publ No. 88. Lyon, International Agency for Research on Cancer, 1987
151. Mustacchi P, Shimkin MB: Cancer of the bladder and infestation with *Schistosoma haematobium*. *J Natl Cancer Inst* 20:825-842, 1958

152. Myslak ZW, Bolt HM, Brockmann W: Tumors of the urinary bladder in painters: A case-control study. *Am J Ind Med* 19:705-713, 1991
153. Najem GR, Louria DB, Seebode JJ, et al: Life time occupation, smoking, caffeine, saccharine, hair dyes and bladder carcinogenesis. *Int J Epidemiol* 11:212-217, 1982
154. National Academy of Sciences: Health Risks of Exposure to Low Levels of Ionizing Radiation (BEIR V). Washington, National Academy Press, 1990
155. National Cancer Institute: Cancer Statistics Review, 1973-1986, Including a Report on the Status of Cancer Control. NIH Publ No. 89-2789. Bethesda, MD, National Cancer Institute, 1989
156. National Research Council: Drinking Water and Health, vol 3. Washington, DC, National Academy Press, 1980, pp 5-21
157. Negri E, Piolatto G, Pira E, et al: Cancer mortality in a northern Italian cohort of rubber workers. *Br J Ind Med* 46:624-628, 1989
158. Nomura A, Kolonel LN, Yoshizawa CN: Smoking, alcohol, occupation and hair dye use in cancer of the lower urinary tract. *Am J Epidemiol* 130:1159-1163, 1989
159. Nomura AMY, Stemmermann GN, Heilbrun LK, et al: Serum vitamin levels and the risk of cancer of specific sites in men of Japanese ancestry in Hawaii. *Cancer Res* 45:2369-2372, 1985
160. Oyasu R, Hirao Y, Izumi K: Enhancement by urine of urinary bladder carcinogenesis. *Cancer Res* 41:478-481, 1981
161. Page WF, Kuntz AJ: Racial and socioeconomic factors in cancer survival. A comparison of Veterans Administration results with selected studies. *Cancer* 45:1029-1040, 1980
162. Parkash O, Kiesswetter H: The role of urine in the etiology of cancer of the urinary bladder. *Urol Int* 31:343-348, 1976
163. Parkes HG, Veys CA, Waterhouse JAH, et al: Cancer mortality in the British rubber industry. *Br J Ind Med* 39:209-220, 1982
164. Payet M: Mortality and morbidity from bladder cancer in West Africa. *Acta Un Internat Contra Cancer* 18:641-642, 1962
165. Pederson-Bjergaard J, Ersboll J, Hansen VL, et al: Carcinoma of the urinary bladder after treatment with cyclophosphamide for non-Hodgkin's lymphoma. *N Engl J Med* 318:1028-1032, 1988
166. Pickle LW, Mason TJ, Howard N, et al: Atlas of US Cancer Mortality among Whites: 1950-80. DHHS Publ No. (NIH) 87-2900. Washington, US Govt Printing Office, 1987
167. Piper JM, Matanoski GM, Tonascia J: Bladder cancer in young women. *Am J Epidemiol* 123:1033-1042, 1986
168. Piper JM, Tonascia J, Matanoski GM: Heavy phenacetin use and bladder cancer in women aged 20 to 49 years. *N Engl J Med* 313:292-295, 1985
169. Puntoni R, Bolognesi C, Bonassi S, et al: Cancer risk evaluation in an area with a high density of chemical plants. An interdisciplinary approach. *Ann NY Acad Sci* 534:808-816, 1988
170. Rebelakos A, Trichopoulos D, Tzonou A, et al: Tobacco smoking, coffee drinking, and occupation as risk factors for bladder cancer in Greece. *J Natl Cancer Inst* 75:455-461, 1985
171. Risch HA, Burch JD, Miller AB, et al: Occupational factors and the incidence of cancer of the bladder in Canada. *Br J Ind Med* 45:361-367, 1988
172. Rockette HE, Arena VC: Mortality studies of aluminum reduction plant workers: Potroom and carbon department. *J Occup Med* 25:549-557, 1983
173. Rowland RG, Henneberry MO, Oyasu R, et al: Effects of urine and continued exposure to carcinogen on progression of early neoplastic urinary bladder lesions. *Cancer Res* 40:4524-4527, 1980
174. Rubino GF, Scansetti G, Piolatto G, et al: The carcinogenic effect of aromatic amines: An epidemiological study on the role of o-toluidine and 4,4'-methylene bis (2-methylaniline) in inducing bladder cancer in man. *Environ Res* 27:241-254, 1982
175. Schairer C, Hartge P, Hoover RN, et al: Racial differences in bladder cancer risk: A case-control study. *Am J Epidemiol* 128:1027-1037, 1988
176. Schiffers E, Jamar J, Renard V: Tobacco and occupation as risk factors in bladder cancer: A case-control study in southern Belgium. *Int J Cancer* 39:287-292, 1987

177. Schulte PA  
cohort expo-  
178. Schulte PA  
Occup Med  
179. Seidman H M  
or dying of can-  
180. Senior Coastguard  
Tuscan Lanning in-  
181. Shaw CH  
presented in  
182. ...

177. Schulte PA, Ringen K, Hemstreet GP, et al: Risk factors for bladder cancer in a cohort exposed to aromatic amines. *Cancer* 58:2156-2162, 1986
178. Schulte PA, Ringen K, Hemstreet GP, et al: Occupational cancer of the urinary tract. *Occup Med: State of the Art Reviews* 2:85-107, 1987
179. Seidman H, Mushinski MH, Gelb SK, et al: Probabilities of eventually developing or dying of cancer—United States, 1985. *CA* 36:36-56, 1985
180. Seniori Costantini A, Paci E, Miligi L, et al: Cancer mortality among workers in the Tuscan tanning industry. *Br J Ind Med* 46:384-388, 1989
181. Shaw GH: Use of Papanicolaou staining to predict urinary tract cancer. Paper presented at the State-of-the-Art Conference on Bladder Cancer Screening. Washington, DC, 1977
182. Silverberg E, Boring CC, Squires TS: Cancer statistics, 1990. *CA* 40:9-26, 1990
183. Silverman DT, Hoover RN, Albert S, et al: Occupation and cancer of the lower urinary tract in Detroit. *J Natl Cancer Inst* 70:237-245, 1983
184. Silverman DT, Hoover RN, Mason TJ, et al: Motor exhaust-related occupations and bladder cancer. *Cancer Res* 46:2113-2116, 1986
185. Silverman DT, Levin LI, Hoover RN: Occupational risks of bladder cancer in the United States: II. Nonwhite men. *J Natl Cancer Inst* 81:1480-1483, 1989
186. Silverman DT, Levin LI, Hoover RN: Occupational risks of bladder cancer among white women in the United States. *Am J Epidemiol* 132:453-461, 1990
187. Silverman DT, Levin LI, Hoover RN, et al: Occupational risks of bladder cancer in the United States: I. White men. *J Natl Cancer Inst* 81:1472-1480, 1989
188. Simon D, Yen S, Cole P: Coffee drinking and cancer of the lower urinary tract. *J Natl Cancer Inst* 54:587-591, 1975
189. Slattery ML, Schumacher MC, West DW, et al: Smoking and bladder cancer: The modifying effect of cigarettes on other factors. *Cancer* 61:402-408, 1988
190. Slattery ML, West DW, Robison LM: Fluid intake and bladder cancer in Utah. *Int J Cancer* 42:17-22, 1988
191. Smith PG, Douglas AJ: Mortality of workers at the Sellafield plant of British Nuclear Fuels. *Br Med J* 293:845-854, 1986
192. Sole G, Sorahan T: Coarse fishing and risk of urothelial cancer. *Lancet* 1:1477-1479, 1985
193. Stasik MJ: Carcinomas of the urinary bladder in a 4-chloro-O-toluidine cohort. *Int Arch Occup Environ Health* 60:21-24, 1988
194. Steenland K, Burnett C, Osorio AM: A case-control study of bladder cancer using city directories as a source of occupational data. *Am J Epidemiol* 126:247-257, 1987
195. Steineck G, Hagman U, Gerhardsson M, et al: Vitamin A supplements, fried foods, fat and urothelial cancer. A case-referent study in Stockholm in 1985-87. *Int J Cancer* 45:1006-1011, 1990
196. Steineck G, Norell SE, Feychting M: Diet, tobacco and urothelial cancer. A 14-year follow-up of 16,477 subjects. *Acta Oncol* 27:323-327, 1988
197. Stern FB, Beaumont JJ, Halperin WE, et al: Mortality of chrome leather tannery workers and chemical exposures in tanneries. *Scand J Work Environ Health* 13:108-117, 1987
198. Tawfik HN: Carcinoma of the urinary bladder associated with schistosomiasis in Egypt: The possible causal relationship. In Miller RW, Watanabe S, Fraumeni JF Jr, et al (eds): *Unusual Occurrences as Clues to Cancer Etiology*. Tokyo, Japan Scientific Societies Press, 1988, pp 197-209
199. Theriault G, DeGuire L, Cordier S: Reducing aluminum: An occupation possibly associated with bladder cancer. *Can Med Assoc J* 124:419-425, 1981
200. Theriault G, Tremblay C, Cordier S, et al: Bladder cancer in the aluminum industry. *Lancet* 1:947-950, 1984
201. Thiede T, Christensen BC: Bladder tumours induced by chlornaphazine. A five-year follow-up study of chlornaphazine-treated patients with polycythaemia. *Acta Med Scand* 185:133-137, 1969
202. Thomas DB, Uhl CN, Hartge P: Bladder cancer and alcoholic beverage consumption. *Am J Epidemiol* 118:720-727, 1983
203. Travis LB, Curtis RE, Boice JD Jr, et al: Bladder cancer after chemotherapy for non-Hodgkin's lymphoma. *N Engl J Med* 321:544-545, 1989

204. Tyler HA, Notley RG, Schweitzer FAW, et al: Vitamin A status and bladder cancer. *Eur J Surg Oncol* 12:35-41, 1986
205. UICC: Screening for Cancer. I. General Principles on Evaluation of Screening for Cancer and Screening for Lung, Bladder and Oral Cancer. Union Internationale Contre le Cancer Technical Report Series, vol 78. Geneva, International Union Against Cancer, 1984
206. US Congress, Office of Technology Assessment: Cancer Testing Technology and Saccharin. Washington, DC, US Govt Printing Office, 1977
207. Vineis P, DiPrima S: Cutting oils and bladder cancer. *Scand J Work Environ Health* 9:449-450, 1983
208. Vineis P, Esteve J, Hartge P, et al: Effects of timing and type of tobacco in cigarette-induced bladder cancer. *Cancer Res* 48:3849-3852, 1988
209. Vineis P, Esteve J, Terracini B: Bladder cancer and smoking in males: Types of cigarettes, age at start, effect of stopping and interaction with occupation. *Int J Cancer* 34:165-170, 1984
210. Vineis P, Magnani C: Occupation and bladder cancer in males: A case-control study. *Int J Cancer* 35:599-606, 1985
211. Wagoner JK: Leukemia and Other Malignancies Following Radiation Therapy for Gynecological Disorders. Unpublished doctoral thesis, Harvard School of Public Health, Boston, 1970
212. Walker AM, Dreyer NA, Friedlander E, et al: An independent analysis of the National Cancer Institute Study on non-nutritive sweeteners and bladder cancer. *Am J Publ Health* 72:376-381, 1982
213. Ward E, Halperin W, Thun M, et al: Bladder tumors in two young males occupationally exposed to MBOCA. *Am J Ind Med* 14:267-272, 1988
214. Wigle DT: Bladder cancer: Possible new high-risk occupation. *Lancet* 2:83-84, 1977
215. Wilkins JR III, Comstock GW: Source of drinking water at home and site-specific cancer incidence in Washington County, Maryland. *Am J Epidemiol* 114:178-190, 1981
216. Williams RR, Horm JW: Association of cancer sites with tobacco and alcohol consumption and socioeconomic status of patients: Interview study from the Third National Cancer Survey. *J Natl Cancer Inst* 58:525-547, 1977
217. Wu M-M, Kuo T-L, Hwang Y-H, et al: Dose-response relation between arsenic concentration in well water and mortality from cancers and vascular diseases. *Am J Epidemiol* 130:1123-1132, 1989
218. Wynder EL, Augustine A, Kabat G, et al: Effect of the type of cigarette smoked on bladder cancer risk. *Cancer* 61:622-627, 1988
219. Wynder EL, Goldsmith R: The epidemiology of bladder cancer: A second look. *Cancer* 40:1246-1268, 1977
220. Wynder EL, Onderdonk J, Mantel N: An epidemiological investigation of cancer of the bladder. *Cancer* 16:1388-1407, 1963
221. Wynder EL, Stellman SD: Comparative epidemiology of tobacco-related cancers. *Cancer Res* 37:4608-4622, 1977
222. Wynder EL, Stellman SD: Artificial sweetener use and bladder cancer: A case-control study. *Science* 207:1214-1216, 1980
223. Yoshida O, Harada T, Miyagawa M, et al: Bladder cancer in workers in the dyeing industry. *Igaku No Ayumi (Japan)* 79:421, 1971
224. Young JL Jr, Percy CL, Asire AJ (eds): Surveillance, Epidemiology, End Results: Incidence and Mortality Data, 1973-77. *Natl Cancer Inst Monogr* 57:1-1082, 1981
225. Zierler S, Feingold L, Danley RA, et al: Bladder cancer in Massachusetts related to chlorinated and chloraminated drinking water: A case-control study. *Arch Environ Health* 43:195-200, 1988

*Address reprint requests to*

Debra T. Silverman, ScD  
Epidemiology and Biostatistics Program  
Division of Cancer Etiology  
National Cancer Institute  
Executive Plaza North, Room 415  
Bethesda, MD 20892